

Table 21.7
Cancer of the Ovary (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2007-2011</u>						
Age at Diagnosis/Death:						
All ages	12.3	13.0	9.8	7.9	8.2	6.6
Under 65	7.5	7.9	5.4	3.1	3.2	2.6
65 and over	45.3	48.0	40.0	41.1	42.7	34.4
All ages (IARC world std) ^c	8.8	9.3	6.8	4.9	5.1	4.2
<u>Age-Specific Rates, 2007-2011</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	0.3	0.2	-	-	-	-
10-14	0.8	0.8	0.7	-	-	-
15-19	1.3	1.4	0.9	0.1	0.1	-
20-24	1.7	1.8	1.4	0.1	0.1	-
25-29	2.3	2.5	2.2	0.3	0.3	0.4
30-34	3.2	3.3	2.7	0.6	0.6	0.6
35-39	5.0	5.2	3.6	1.1	1.1	1.1
40-44	9.0	9.5	5.8	2.3	2.3	2.2
45-49	14.3	15.0	9.4	4.7	5.0	4.0
50-54	20.4	21.4	13.5	8.8	9.2	7.1
55-59	25.5	27.3	18.0	13.7	14.3	11.2
60-64	31.4	33.4	24.9	20.1	21.0	16.7
65-69	38.0	40.8	33.1	28.3	29.5	24.9
70-74	45.2	48.4	39.2	36.9	38.6	32.4
75-79	48.0	50.8	40.9	44.9	46.7	37.5
80-84	50.8	53.4	47.1	53.9	55.8	44.4
85+	50.3	52.2	47.0	56.4	58.3	42.3

- ^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.
- ^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.
- ^c Rates are per 100,000 and are age-adjusted to the IARC world standard population.
- Statistic not shown. Rate based on less than 16 cases for the time interval.